

Effect of Pneumococcal Vaccination on Nasopharyngeal Carriage of *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, and *Staphylococcus aureus* in Fijian Children

Eileen M. Dunne,^a Jayne Manning,^a Fiona M. Russell,^a Roy M. Robins-Browne,^{b,c} E. Kim Mulholland,^{a,d,e} and Catherine Satzke^{a,b,c}

Centre for International Child Health^a and Infectious Diseases & Microbiology,^b Murdoch Childrens Research Institute, Royal Children's Hospital, Parkville, VIC, Australia; Department of Microbiology & Immunology, The University of Melbourne, Parkville, VIC, Australia^c; London School of Hygiene & Tropical Medicine, London, United Kingdom^d; and Menzies School of Health Research, Darwin, NT, Australia^e

The 7-valent pneumococcal conjugate vaccine (PCV7) reduces carriage of vaccine type *Streptococcus pneumoniae* but leads to replacement by nonvaccine serotypes and may affect carriage of other respiratory pathogens. We investigated nasopharyngeal carriage of *S. pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, and *Staphylococcus aureus* in Fijian infants participating in a pneumococcal vaccine trial using quantitative PCR. Vaccination did not affect pathogen carriage rates or densities, whereas significant differences between the two major ethnic groups were observed.

Nasopharyngeal (NP) carriage of pathogenic bacteria is the primary reservoir for maintaining bacterial species within a population (7) and is considered a prerequisite for development of major childhood diseases, including bacterial pneumonia, meningitis, and otitis media. *Streptococcus pneumoniae* is the most common bacterial cause of childhood pneumonia and is responsible for at least 800,000 child deaths annually, primarily in developing countries (18, 33). While rarely fatal, otitis media is the most frequently reported childhood bacterial infection: approximately 80% of children experience otitis media by age three (17). *S. pneumoniae*, *Haemophilus influenzae*, and *Moraxella catarrhalis* are the predominant causes of otitis media (36).

PCV7 (Prevnar; Pfizer Inc.) effectively reduces carriage and subsequent disease caused by the serotypes of *S. pneumoniae* included in the vaccine. However, pneumococcal vaccination has minimal impact on the overall rate of pneumococcal carriage due to replacement by nonvaccine serotypes (3, 8, 11). Reports demonstrating an inverse relationship between nasopharyngeal carriage of vaccine type *S. pneumoniae* and *Staphylococcus aureus* (2, 21, 22) and increases in the proportion of otitis media caused by nontypeable *H. influenzae* following PCV7 vaccination (1, 4, 41) generated concern that removal of vaccine type pneumococci from the nasopharynx could facilitate colonization by other respiratory pathogens. This study aimed to evaluate the effects of pneumococcal immunization on carriage of respiratory pathogens in a high-risk population.

A quantitative real-time PCR (qPCR) method was developed to measure *S. pneumoniae*, *H. influenzae*, *M. catarrhalis*, and *S. aureus* and applied to nasopharyngeal swabs collected from 17-month-old participants in a phase II pneumococcal vaccine trial in Suva, Fiji. The trial investigated appropriate infant dosing strategy of PCV7 and effects of a 23-valent polysaccharide (23vPPS; Pneumovax; Merck & Co., Inc.) booster given at 12 months. Enrollment, ethical approval, and swab collection and storage have been detailed previously (25, 27, 29). Swabs (collected in 2006 and 2007) in STGG media (19) were transported to Melbourne, Australia, on dry ice and stored at -80°C until use. The following sample groups were examined: group A ($n = 54$), who received three doses of PCV7 at 6, 10, and 14 weeks of age; group B ($n = 48$), who received three PCV7 doses plus a booster of 23vPPS at 12

months of age; and group H ($n = 59$), who were unvaccinated. Previous examination of pneumococcal serotypes showed that three doses of PCV7 led to a reduction in carriage of vaccine type *S. pneumoniae* that was sustained to the age of 17 months, whereas carriage rates of non-PCV serotypes were similar, and that 23vPPS had no impact on carriage (27).

Cells from a 100- μl aliquot of sample were lysed in 200 μl of 50 mM phosphate buffer, pH 6.7, containing 1 mg/ml lysozyme, 0.075 mg/ml mutanolysin, and 2 mg/ml proteinase K and incubated at 56°C for 45 min. An additional 2 mg/ml proteinase K and 1% SDS were added, and then cells were incubated at 56°C for 10 min. DNA was extracted and purified into 100 μl elution buffer using the QIAmp DNA minikit (Qiagen). DNA for standard curves was extracted from *S. pneumoniae* ATCC 6305, *H. influenzae* F412 (kindly provided by Cynthia Whitchurch, University of Technology, Sydney, Australia), *S. aureus* ATCC 29213, and *M. catarrhalis* ATCC 8176. We developed two quantitative duplex PCR assays, one to detect *S. pneumoniae* and *S. aureus* and another to detect *H. influenzae* and *M. catarrhalis*. *S. pneumoniae* was detected using previously published primers and probes (30). Primers for *S. aureus* detection were modified for use with a previously published probe (6) for enhanced sensitivity. Sequence alignment of *M. catarrhalis copB* from 14 available GenBank entries revealed that published probes (9, 30) hybridize to a variable region of the gene, so a new primer/probe set was designed to target a conserved region of this gene. This qPCR was coupled with an assay for *H. influenzae* that was developed to detect both typeable and nontypeable *H. influenzae* but not the closely related *Haemophilus haemolyticus* (38). Two microliters of DNA was used in each of two duplex qPCRs performed with Brilliant III Ultra-Fast QPCR master mix (Agilent Technologies) on a Stratagene Mx3005 real-time PCR instrument with an initial activation of 95°C for 3 min

Received 4 December 2011 Accepted 7 December 2011

Published ahead of print 14 December 2011

Address correspondence to Eileen M. Dunne, eileen.dunne@mcri.edu.au.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JCM.06589-11

TABLE 1 Sequences and concentrations of primers and probes for duplex quantitative PCRs to detect *S. pneumoniae* and *S. aureus* (reaction 1) and *M. catarrhalis* and *H. influenzae* (reaction 2)

Reaction	Species and target gene	Primer and probe sequences ^a (5'-3') [concn (nM)]	Reference and/or source
1	<i>S. pneumoniae</i> <i>lytA</i>	F, TCTTACGCAATCTAGCAGATGAAGC [96]; R, GTTGTTGGTTGGTTATTCGTGC [96]; probe, FAM-TTTGCCGAAAACGCTTGATACAGGG-BHQ1 [96]	30
	<i>S. aureus</i> <i>nuc</i>	F, GTTGCTTAGTGTAACTTTAGTTGTA [288]; R, GCACTATATACTGTTGGATCTTCAGAA [288]; probe, TxR-TGCATCACAAACAGATAACGGCGTAAATAGAAG-BHQ2 [192]	6, this study
2	<i>M. catarrhalis</i> <i>copB</i>	F, CGTGTGACCGTTTGGACTTT [96]; R, TAGATTAGGTTACCGCTGACG [96]; probe, Cy5-ACCGACATCAACCCAAGCTTTGG-BHQ3 [96]	This study
	<i>H. influenzae</i> <i>hpd</i>	F, GGTAAATATGCCGATGGTGTG [96]; R, TGCATCTTTACGCACGGTGTA [288]; probe, HEX-TTGTGTACACTCCGT“T”GGTAAAAGAACTGACAC-SpacerC3 ^b [96]	38

^a F, forward; R, reverse; FAM, 6-carboxyfluorescein; BHQ1, black hole quencher 1; TxR, Texas red; HEX, hexachloro-6-carboxyfluorescein.

^b “T”, BHQ-1 (dT).

followed by 40 cycles of 95°C for 20 s and 60°C for 20 s. Table 1 details PCR primers (Sigma-Aldrich) and dually labeled probes (Eurogentec).

qPCR data were used to determine the carriage rate (% positive) and carriage density (CFU/ml) for each bacterial species. Fisher's exact test and the Mann-Whitney test were used to evaluate differences in rate and density, respectively. *P* values <0.05 were considered statistically significant. Associations were measured by calculating odds ratios (OR) and 95% confidence intervals (CI 95%).

Of the 161 NP swabs examined, at least one pathogen was identified in 141 (87.5%). *M. catarrhalis* was identified most frequently, found in 122 (75.8%) swabs, followed by *S. pneumoniae* (*n* = 92; 57.1%) and *H. influenzae* (*n* = 72; 44.7%). The overall carriage rate of *S. aureus*, 3.3%, was consistent with reports from the Gambia (12) and Western Australia, Australia (39). *S. aureus* carriage is typically highest in neonates and older children (2, 22, 23), and the nose, rather than the nasopharynx, is the primary ecological niche (40). Due to its low frequency, *S. aureus* was not included in subsequent analyses. The higher colonization rates of *S. pneumoniae*, *H. influenzae*, and *M. catarrhalis* were similar to those observed in high-risk populations (12, 16, 30, 39). Cocolonization was common, with 97 of 161 (60.2%) children colonized by multiple species compared with 44 (27.3%) colonized by one of the species tested. Positive associations between colonization by *S. pneumoniae* and *M. catarrhalis* (OR, 3.16; CI 95%, 1.49 to 6.71), *S. pneumoniae* and *H. influenzae* (OR, 3.85; CI 95%, 1.95 to 7.58), and *M. catarrhalis* and *H. influenzae* (OR, 6.70; CI 95%, 2.70 to 17.60) were found.

Regardless of 23vPPS booster status, vaccination with PCV7 did not affect carriage rates (Table 2) or densities (data not shown) of the four pathogens examined. The population of Fiji consists of

57% indigenous Fijians and 38% Indo-Fijians of Indian descent. The original vaccine trial was designed to approximate this breakdown in each vaccine group. As indigenous Fijian children are known to have higher rates of *S. pneumoniae* carriage (26), data were stratified by ethnicity and reanalyzed. Swabs from groups A, B, and H were collected from 99 Fijians and 54 Indo-Fijians (ethnicity self-reported upon enrollment). Indigenous Fijians (i) were more likely to carry two or more organisms (OR, 9.66; CI 95%, 4.49 to 20.77), (ii) had significantly higher carriage rates of *S. pneumoniae*, *H. influenzae*, and *M. catarrhalis* (Table 2), and (iii) had higher carriage densities of *S. pneumoniae* and *M. catarrhalis* (Fig. 1) than Indo-Fijians.

Next, we examined the effects of pneumococcal vaccination separately for each ethnic group. No differences in carriage rates were observed (data not shown). Vaccination did not affect carriage densities of *S. pneumoniae* or *H. influenzae* in either indigenous Fijians or Indo-Fijians (Fig. 2A and B). Carriage densities of *M. catarrhalis* were significantly higher in indigenous Fijian children who received three doses of PCV7 plus a 23vPPV booster than in those who were unvaccinated or received PCV7 alone (Fig. 2C). This increase in *M. catarrhalis* density was not observed in Indo-Fijian children.

In summary, vaccination of Fijian infants with PCV7 alone or with the 23vPPS booster did not affect nasopharyngeal carriage rates or densities of *S. pneumoniae*, *H. influenzae*, *M. catarrhalis*, and *S. aureus*. However, significant differences between the two major ethnic groups were found, with indigenous Fijians more likely to carry higher densities of multiple respiratory pathogens than Fijians of Indian descent.

Several studies investigated effects of pneumococcal vaccination on *S. aureus*, and most did not find any increases in *S. aureus* colonization associated with PCV7 (4, 5, 13) or PCV9 (14) vac-

TABLE 2 Nasopharyngeal carriage rates of *S. pneumoniae*, *H. influenzae*, and *M. catarrhalis* in Fijian children stratified by vaccine group and ethnicity

Species	No. of positive samples/total no. of samples ^a (%) for:				
	Vaccine group			Fijian group	Indo-Fijian group
	3 doses PCV7	3 doses PCV7 + 23vPPV	No vaccine		
<i>S. pneumoniae</i>	30/54 (56)	29/48 (60)	33/59 (56)	68/99 (69)	20/54 (37)*
<i>H. influenzae</i>	24/54 (44)	24/48 (50)	24/59 (41)	66/99 (67)	6/54 (11)*
<i>M. catarrhalis</i>	42/54 (78)	39/48 (81)	41/59 (70)	91/99 (92)	24/54 (44)*

^a *, *P* < 0.001. For all other comparisons, *P* was >0.05.




FIG 1 Nasopharyngeal carriage densities of *S. pneumoniae* (*S. pneu*), *H. influenzae* (*H. inf*), and *M. catarrhalis* (*M. cat*) in indigenous Fijian (Fijian) and Indo-Fijian infants. The median and interquartile ranges are shown in gray. *, $P < 0.05$. For all other comparisons, P was >0.05 .

nation. In contrast, van Gils et al. reported a temporary increase in *S. aureus* colonization at 12 months of age in infants who received three doses of PCV7 (34). The low carriage rate of *S. aureus* in our study hindered our ability to assess the impact of pneumococcal vaccination on its carriage in Fijian infants. Reported increases in the proportion of otitis media caused by *H. influenzae* and *M. catarrhalis* associated with PCV7 vaccination (1, 4, 41) may reflect

the reduction in cases caused by vaccine type *S. pneumoniae* rather than true increases in disease caused by other pathogens. For example, Stamboulidis et al. noted that, following PCV7 immunization, the overall rate of acute otitis media with otorrhea and the incidence of *S. pneumoniae*- and *H. influenzae*-caused otorrhea declined, yet *H. influenzae* replaced *S. pneumoniae* as the predominant organism (31).

Revai et al. examined nasopharyngeal colonization during acute otitis media and found an increase in *M. catarrhalis* in PCV7-vaccinated children compared with unvaccinated historical controls (24). Consistent with our findings, studies on the effects of pneumococcal vaccination in healthy children found no increases in carriage of *H. influenzae* (14, 35) or *M. catarrhalis* (35). We used the final swabs collected during the study, taken at 17 months, 14 months after the last dose of PCV7 and 5 months after the 23vPPS booster. Therefore, any transient effects of pneumococcal vaccination would not have been observed in this study.

We found strong positive associations between carriage of *S. pneumoniae*, *H. influenzae*, and *M. catarrhalis*. Other researchers have reported similar associations between *S. pneumoniae* and *H. influenzae* (10, 12, 14), *S. pneumoniae* and *M. catarrhalis* (10, 12), and *M. catarrhalis* and *H. influenzae* (10, 37) in children of a comparable age. Pettigrew et al. (20) found negative associations between *S. pneumoniae* and *H. influenzae* and between *M. catarrhalis* and *H. influenzae* in children with upper respiratory infections, suggesting that the presence of an active infection may influence colonization dynamics. In our study, attributes of the host rather




FIG 2 Nasopharyngeal carriage densities of *S. pneumoniae* (A), *H. influenzae* (B), and *M. catarrhalis* (C) separated by vaccine group and ethnicity. The median and interquartile ranges are shown in gray. *, $P < 0.01$. For all other comparisons, P was >0.05 .

than the presence of a particular bacterial species seemed to be the dominant factor relating to the cocarriage of multiple pathogens. Host immunity likely plays a role in the negative association between *S. aureus* and *S. pneumoniae*, as this inverse relationship was not observed in HIV-positive children (14).

Indigenous Fijians have been shown to have higher incidence of pneumonia (15), invasive pneumococcal disease (28), and group A streptococcal disease (32) than Indo-Fijians. Reasons for this are unclear but may include genetic susceptibility, cultural practices, and/or environmental factors such as crowded living conditions. In our study, antibiotic use and monthly annual income levels were similar between the ethnicities but Fijian families had more children under 5 in the household, a risk factor for *S. pneumoniae* colonization (26). Differences in carriage between ethnic groups raise the possibility that vaccines targeting colonizing bacteria may have differential effects in different populations. Indeed, the 23vPPS booster was associated with increased carriage densities of *M. catarrhalis* in indigenous Fijians but not Indo-Fijians. Differences in susceptibilities to bacterial colonization warrant further investigation, and the high levels of carriage of multiple pathogens common in developing countries should be taken into consideration when designing and monitoring new pneumococcal vaccine introduction.

ACKNOWLEDGMENTS

This work was supported by funding from the Murdoch Childrens Research Institute and the Victorian Government's Operational Infrastructure Support Program.

We thank Heidi Smith-Vaughan and Michael Binks from the Menzies School of Health Research and Xin Wang from the Centers for Disease Control and Prevention for advice on assay development. We acknowledge the PneuCarriage Project team for expert technical assistance, the MCRI Clinical Epidemiology and Biostatistics Unit for helpful advice, and Eleanor Neal for manuscript editing. We sincerely thank all the families and staff on the Fiji Pneumococcal project.

REFERENCES

- Block SL, et al. 2004. Community-wide vaccination with the heptavalent pneumococcal conjugate significantly alters the microbiology of acute otitis media. *Pediatr. Infect. Dis. J.* 23:829–833.
- Bogaert D, et al. 2004. Colonisation by *Streptococcus pneumoniae* and *Staphylococcus aureus* in healthy children. *Lancet* 363:1871–1872.
- Brugger SD, Frey P, Aebi S, Hinds J, Mühlemann K. 2010. Multiple colonization with *S. pneumoniae* before and after introduction of the seven-valent conjugated pneumococcal polysaccharide vaccine. *PLoS One* 5:e11638.
- Casey JR, Adlowitz DG, Pichichero ME. 2010. New patterns in the otopathogens causing acute otitis media six to eight years after introduction of pneumococcal conjugate vaccine. *Pediatr. Infect. Dis. J.* 29:304–309.
- Cohen R, et al. 2007. Pneumococcal conjugate vaccine does not influence *Staphylococcus aureus* carriage in young children with acute otitis media. *Clin. Infect. Dis.* 45:1583–1587.
- Elizaquível P, Aznar R. 2008. A multiplex RTi-PCR reaction for simultaneous detection of *Escherichia coli* O157:H7, *Salmonella* spp. and *Staphylococcus aureus* on fresh, minimally processed vegetables. *Food Microbiol.* 25:705–713.
- García-Rodríguez JÁ, Fresnadillo Martínez MJ. 2002. Dynamics of nasopharyngeal colonization by potential respiratory pathogens. *J. Antimicrob. Chemother.* 50:59–74.
- Ghaffar F, et al. 2004. Effect of the 7-valent pneumococcal conjugate vaccine on nasopharyngeal colonization by *Streptococcus pneumoniae* in the first 2 years of life. *Clin. Infect. Dis.* 39:930–938.
- Greiner O, Day PJ, Altwegg M, Nadal D. 2003. Quantitative detection of *Moraxella catarrhalis* in nasopharyngeal secretions by real-time PCR. *J. Clin. Microbiol.* 41:1386–1390.
- Jacoby P, et al. 2007. Modelling the co-occurrence of *Streptococcus pneumoniae* with other bacterial and viral pathogens in the upper respiratory tract. *Vaccine* 25:2458–2464.
- Kellner JD, et al. 2008. Effects of routine infant vaccination with the 7-valent pneumococcal conjugate vaccine on nasopharyngeal colonization with *Streptococcus pneumoniae* in children in Calgary, Canada. *Pediatr. Infect. Dis. J.* 27:526–532.
- Kwambana B, Barer MR, Bottomley C, Adegbola RA, Antonio M. 2011. Early acquisition and high nasopharyngeal co-colonisation by *Streptococcus pneumoniae* and three respiratory pathogens amongst Gambian newborns and infants. *BMC Infect. Dis.* 11:175.
- Lee GM, et al. 2009. Epidemiology and risk factors for *Staphylococcus aureus* colonization in children in the post-PCV7 era. *BMC Infect. Dis.* 9:110.
- Madhi SA, et al. 2007. Long-term effect of pneumococcal conjugate vaccine on nasopharyngeal colonization by *Streptococcus pneumoniae*—and associated interactions with *Staphylococcus aureus* and *Haemophilus influenzae* colonization—in HIV-infected and HIV-uninfected children. *J. Infect. Dis.* 196:1662–1666.
- Magree HC, et al. 2005. Chest X-ray-confirmed pneumonia in children in Fiji. *Bull. World Health Organ.* 83:427–433.
- Montgomery JM, et al. 1990. Bacterial colonization of the upper respiratory tract and its association with acute lower respiratory tract infections in highland children of Papua New Guinea. *Rev. Infect. Dis.* 12:S1006–S1016.
- Murphy TF, Parameswaran GI. 2009. *Moraxella catarrhalis*, a human respiratory tract pathogen. *Clin. Infect. Dis.* 49:124–131.
- O'Brien KL, et al. 2009. Burden of disease caused by *Streptococcus pneumoniae* in children younger than 5 years: global estimates. *Lancet* 374:893–902.
- O'Brien KL, et al. 2001. Evaluation of a medium (STGG) for transport and optimal recovery of *Streptococcus pneumoniae* from nasopharyngeal secretions collected during field studies. *J. Clin. Microbiol.* 39:1021–1024.
- Pettigrew MM, Gent JF, Revai K, Patel JA, Chonmaitree T. 2008. Microbial interactions during upper respiratory tract infections. *Emerg. Infect. Dis.* 14:1584–1591.
- Quintero B, et al. 2011. Epidemiology of *Streptococcus pneumoniae* and *Staphylococcus aureus* colonization in healthy Venezuelan children. *Eur. J. Clin. Microbiol. Infect. Dis.* 30:7–19.
- Regev-Yochay G, et al. 2004. Association between carriage of *Streptococcus pneumoniae* and *Staphylococcus aureus* in children. *JAMA* 292:716–720.
- Regev-Yochay G, et al. 2009. Parental *Staphylococcus aureus* carriage is associated with staphylococcal carriage in young children. *Pediatr. Infect. Dis. J.* 28:960–965.
- Revai K, et al. 2006. Effect of pneumococcal conjugate vaccine on nasopharyngeal bacterial colonization during acute otitis media. *Pediatrics* 117:1823–1829.
- Russell FM, et al. 2009. Immunogenicity following one, two, or three doses of the 7-valent pneumococcal conjugate vaccine. *Vaccine* 27:5685–5691.
- Russell FM, et al. 2006. Pneumococcal nasopharyngeal carriage and patterns of penicillin resistance in young children in Fiji. *Ann. Trop. Paediatr.* 26:187–197.
- Russell FM, et al. 2010. Pneumococcal nasopharyngeal carriage following reduced doses of a 7-valent pneumococcal conjugate vaccine and a 23-valent pneumococcal polysaccharide vaccine booster. *Clin. Vaccine Immunol.* 17:1970–1976.
- Russell FM, et al. 2010. Invasive pneumococcal disease in Fiji: clinical syndromes, epidemiology, and the potential impact of pneumococcal conjugate vaccine. *Pediatr. Infect. Dis. J.* 29:870–872.
- Russell FM, et al. 2010. Safety and immunogenicity of the 23-valent pneumococcal polysaccharide vaccine at 12 months of age, following one, two, or three doses of the 7-valent pneumococcal conjugate vaccine in infancy. *Vaccine* 28:3086–3094.
- Smith-Vaughan H, et al. 2006. Measuring nasal bacterial load and its association with otitis media. *BMC Ear Nose Throat Disord.* 6:10.
- Stamboulidis K, et al. 2011. The impact of the heptavalent pneumococcal conjugate vaccine on the epidemiology of acute otitis media complicated by otorrhea. *Pediatr. Infect. Dis. J.* 30:551–555.
- Steer AC, et al. 2008. High burden of invasive beta-haemolytic streptococcal infections in Fiji. *Epidemiol. Infect.* 136:621–627.

33. van der Poll T, Opal SM. 2009. Pathogenesis, treatment, and prevention of pneumococcal pneumonia. *Lancet* 374:1543–1556.
34. van Gils EJM, et al. 2011. Effect of seven-valent pneumococcal conjugate vaccine on *Staphylococcus aureus* colonisation in a randomised controlled trial. *PLoS One* 6:e20229.
35. van Gils EJM, Veenhoven RH, Rodenburg GD, Hak E, Sanders EAM. 2011. Effect of 7-valent pneumococcal conjugate vaccine on nasopharyngeal carriage with *Haemophilus influenzae* and *Moraxella catarrhalis* in a randomized controlled trial. *Vaccine* 29:7595–7598.
36. Vergison A. 2008. Microbiology of otitis media: a moving target. *Vaccine* 26:G5–G10.
37. Verhaegh SJC, et al. 2011. Colonization of healthy children by *Moraxella catarrhalis* is characterized by genotype heterogeneity, virulence gene diversity and co-colonization with *Haemophilus influenzae*. *Microbiology* 157:169–178.
38. Wang X, et al. 2011. Detection of bacterial pathogens in Mongolia meningitis surveillance with a new real-time PCR assay to detect *Haemophilus influenzae*. *Int. J. Med. Microbiol.* 301:303–309.
39. Watson K, et al. 2006. Upper respiratory tract bacterial carriage in Aboriginal and non-Aboriginal children in a semi-arid area of Western Australia. *Pediatr. Infect. Dis. J.* 25:782–790.
40. Wertheim HFL, et al. 2005. The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infect. Dis.* 5:751–762.
41. Wiertsema SP, et al. 2011. Predominance of nontypeable *Haemophilus influenzae* in children with otitis media following introduction of a 3 + 0 pneumococcal conjugate vaccine schedule. *Vaccine* 29:5163–5170.